Name:

1 Anti Derivative and u sub

- 1. Compute the following Integrals
 - (a) $\int \sqrt{3x} + 4 \sec(x) \tan(x) \frac{2}{\sqrt{1 x^2}} dx$ (b) $\int \frac{3x^2 - 4\sqrt{x} + 3}{x} dx$ (c) $\int \csc(x) [\cot(x) + \sin(x)] dx$ (d) $\int \frac{x}{1 + x^2} dx$ (e) $\int \frac{x}{1 + x^4} dx$ (f) $\int \frac{e^x}{1 + e^{2x}} dx$ (g) $\int \sin^3(2x) \cos(2x) dx$ (h) $\int \sin(3x) \cos^{1/2}(3x) dx$ (i) $\int \sin(x) \sec^2(\cos(x)) dx$

2 Definition of the integral

- 2. Let f(x) = 3x + 4 over the interval [1, 6]. Let n = 5.
 - (a) Graph the function with n regions. Label the important points on the graph.
 - (b) Compute the LH rule Riemann sum.
 - (c) Compute the RH rule Riemann sum.

3. Using the **definition** of the integral compute

$$\int_{1}^{4} 3x - 2\,dx.$$

4. Using the **FTC II** compute

$$\int_{1}^{4} 3x - 2\,dx.$$

5. Using the **FTC I** compute

(a)
$$\frac{d}{dx} \left[\int_{4}^{x^2} f(t) dt \right]$$

(b) $\frac{d}{dx} \left[\int_{4}^{3x+2} e^{t^3} dt \right]$
(c) $\frac{d}{dx} \left[\int_{x}^{2x} f(t) dt \right]$

- 6. For $f(x) = x^3$ and [0,3] find f_{ave} . And find the *c* from the MVTI.
- 7. For $f(x) = 1 + x^2$ and [-2, 0] find f_{ave} . And find the *c* from the MVTI.

3 Application of the integral

- 8. Velocity, Acceleration and Position
 - (a) Let a(t) = -5sin(t), v(0) = 5 and s(0) = -7.
 - i. Find v(t) and s(t).
 - ii. When does the object stop?
 - (b) Let a(t) = -12t, v(0) = 6 and s(0) = 0.
 - i. Find v(t) and s(t).
 - ii. When does the object stop?
 - iii. What is the position of the object when it stops?
- 9. Find the area between the functions $y = x^2$ and y = 4.
- 10. Find the area between the functions $y = x^2$ and y = x + 1.
- 11. Find the area between the functions $y = e^{3x}$, y = 4 and the y-axis.

- 12. Find the area between the functions $x = y^2$ and x = 4.
- 13. Find the area between the functions $x = y^2$ and y = x 1.
- 14. Find the area between the functions $y = \ln(x)$, x = 1 and y = 4.
- 15. Find the area between the functions $y = \ln(x)$, x = 3 and the x-axis.
- 16. Find the volume of the solid formed when rotating the region bounded by $y = x^2$ and y = 4 around the x-axis using discs.
- 17. Find the volume of the solid formed when rotating the region bounded by $y = x^2$ and y = 4 in the first quadrant around the y-axis using discs.
- 18. Find the volume of the solid formed when rotating the region bounded by $y = e^{3x}$, y = 4 and the y-axis around the x-axis using discs.
- 19. Find the volume of the solid formed when rotating the region bounded by $y = x^2$ and y = 4 around the x-axis using shells.
- 20. Find the volume of the solid formed when rotating the region bounded by $y = x^2$ and y = 4x around the y-axis using shells.
- 21. Rotate the region bounded by y = 3x, y = 4 and the y-axis around the y-axis using discs.
 - (a) Set up the integral with discs and with shells.
 - (b) Compute both of the integrals and compare.
- 22. Rotate the region bounded by $y = e^{3x}$, y = 4 and the y-axis around the y-axis using discs.
 - (a) Set up the integral with discs and with shells.
 - (b) Compute one of the integrals.
- 23. Rotate the region bounded by the ellipse $x^2 + \frac{y^2}{4} = 1$, y = 2x 2 around the x-axis using discs.
 - (a) Set up the integral with discs and with shells.
 - (b) Compute both of the integrals and compare.
- 24. Compute the following integrals using by parts.
 - (a) $\int x e^{2x} dx$

- (b) $\int x^2 e^{3x} dx$
- (c) $\int x^3 e^{3x^2} dx$
- (d) $\int x \sin(2x) dx$
- (e) $\int x^2 \sin(2x) dx$
- (f) $\int \ln(x) dx$
- (g) $\int x \ln(x) dx$
- (h) $\int x^3 \ln(x) dx$
- (i) $\int \arctan(2x) dx$

25. Compute the following integrals using u-sub.

- (a) $\int xe^{x^{2}} dx$ (b) $\int x \sec^{2}(x^{2}) dx$ (c) $\int (x^{2}+2)e^{x^{3}+6x} dx$ (d) $\int x\sqrt{x-1} dx$ (e) $\int x\sqrt{x^{2}-1} dx$ (f) $\int \frac{3x}{1+x^{2}} dx$ (g) $\int \frac{4}{1+x^{2}} dx$ (h) $\int \sec^{2}(x) \sin(\tan(x)) dx$ (i) $\int e^{2x} \sec(e^{2x}) \tan(e^{2x}) dx$ (j) $\int \tan(x) dx$. Hint use $\tan(x) = \frac{\sin(x)}{\cos(x)}$ and u-sub.
 - (k) $\int \cot(x) dx$
 - (l) $\int \frac{e^x}{1+e^{2x}} dx$
- (m) $\int \frac{e^{2x}}{1+e^{2x}} dx$