## Math 3520 - Test 2 Review

## 1 Number Theory

- 1. Let  $a, b, c, d \in \mathbb{Z}$  with  $b \neq 0$ . Show if a|b and b|c then a|c.
- 2. Let  $a, b, c, x, y \in \mathbb{Z}$ . Show if a|b and a|c then a|ax + by.
- 3. Let  $a, b, c, d \in \mathbb{Z}$  with  $c \neq 0$ . Show if a|b and c|d then ac|ad + bc.
- 4. Prove  $3|n^3 n$  for every integer n.
- 5. Let  $a, b, c \in \mathbb{Z}$ . If a | bc and gcd(a, b) = 1 then a | c.
- 6. Let  $a, b \in \mathbb{Z}$  and let p be a prime. If p|ab then p|a or p|b.
- 7. Let  $a, b, c \in \mathbb{Z}$  where gcd(a, b) = 1. If a|c and b|c then ab|c.
- 8. Let  $n \in \mathbb{Z}$  be odd. Prove  $n^2 \equiv 1 \mod 4$ .
- 9. Let  $n \in \mathbb{Z}$  be odd. Prove  $n^2 \equiv 1 \mod 8$ .
- 10. Let  $a, b, c \in \mathbb{Z}$  so that  $a^2 + b^2 = c^2$ . Prove 4|ab (hint use 9).
- 11. Prove or disprove if  $a, b \in \mathbb{Z}$  and a, b are odd then  $8|a^2 b^2$ .
- 12. Prove  $\sqrt{3}$  is irrational.
- 13. For the following pairs of numbers, find their gcd and find a linear combination of the numbers equal to their gcd.
  - (a) a = 78 and b = 48
  - (b) a = 79 and b = 49
  - (c) a = 253 and b = 207

## 2 Group Theory

- 14. State the definition of an Algebraic structure and a binary operation.
- 15. State the definition of a group.
- 16. State the definition of an Abelian group.

- 17. State and prove the cancellation law.
- 18. Are the following algebraic structures groups? If it is a group prove it. If not prove it is not a group.
  - (a)  $(\mathbb{Q}^*, \cdot)$  where  $\cdot$  is regular multiplication.
  - (b)  $(\mathbb{Z}, \odot)$  where  $a \odot b = a + b 2$ .
  - (c)  $(\mathbb{Z}, \circledast)$  where  $a \circledast b = ab + a + b$ .
  - (d)  $(\mathbb{Q} \setminus \{-1\}, \circledast)$  where  $a \circledast b = ab + a + b$ .
- 19. For the following algebraic structures write the operation tables. State if the structure is a group or not. If not state which property fails and how. Identify the identity in each case.
  - (a)  $(\mathbb{Z}_5, +)$
  - (b)  $(\mathbb{Z}_5, \cdot)$
  - (c)  $(\mathbb{Z}_{6}^{*}, \cdot)$
  - (d)  $(S_3, \circ)$
  - (e)  $(D, \circ)$  where

$$D = \left\{ \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 1 & 2 & 3 \end{array} \right), \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 1 & 3 & 2 \end{array} \right), \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 2 & 1 & 3 \end{array} \right), \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 3 & 1 & 2 \end{array} \right) \right\}$$

(f)  $(D, \circ)$  where  $D \subseteq S_3$  defined as

$$D = \left\{ \left( \begin{array}{rrr} 1 & 2 & 3 \\ 1 & 2 & 3 \end{array} \right), \left( \begin{array}{rrr} 1 & 2 & 3 \\ 2 & 3 & 1 \end{array} \right), \left( \begin{array}{rrr} 1 & 2 & 3 \\ 3 & 1 & 2 \end{array} \right) \right\}$$

(g)  $(M, \cdot)$  where  $M \subseteq GL_2(\mathbb{R})$  defined as

$$M = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right\}$$

- 20. We showed in class that some groups are "the same" by finding a function from one group to the other that preserved the operation. The following pairs of groups are "the same" in that sense find the function and verify that it preserves the operation.
  - (a)  $(Z_4, +)$  and  $(Z_5^*, \cdot)$

(b)  $(Z_2, +)$  and  $(M, \cdot)$  where  $M \subseteq GL_2(\mathbb{R})$  defined as

$$M = \left\{ \left[ \begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right], \left[ \begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array} \right] \right\}$$

(c)  $(Z_{12}^*, \cdot)$  and  $(\mathbb{Z}_2 \times \mathbb{Z}_2, +)$ 

- 21. Let G be a group prove
  - (a) The identity in G is unique.
  - (b) If  $a \in G$  then  $a^{-1}$  is unique.
  - (c) If  $a \in G$  then  $(a^{-1})^{-1} = a$ .
  - (d) If  $a, b \in G$  then  $(ab)^{-1} = b^{-1}a^{-1}$ .
- 22. Let G be a group prove and let  $a, b \in G$ . If a and b commute then  $a^{-1}$  and  $b^{-1}$  commute.
- 23. Let (G, \*) be a group prove and let  $a, b, c \in G$ . Prove the following equations have an unique solution  $x \in G$ .
  - a \* x \* c = b
  - c \* a \* x = b
- 24. Let (G, \*) be a group prove. Assume there is some element  $f \in G$  so that for some element  $a \in G$ we have af = a. Is f the identity? Prove or disprove.
- 25. Let (G, \*) be a group prove. If a \* a = e for all  $a \in G$  then G is Abelian.

## 3 Sub Groups

- 26. Definition of a **subgroup** and of an **isomorphism**.
- 27. For the group  $(\mathbb{Z}, +)$  prove, using the 2-step subgroup test, that  $H = \{3n : n \in \mathbb{Z}\}$  is a subgroup.
- 28. For the group  $(\mathbb{Z}, +)$  Show  $H = \{2n + 1 : n \in \mathbb{Z}\}$  is not a subgroup.

29. For the group  $(S_3, \circ)$  Show

$$H = \left\{ \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 1 & 2 & 3 \end{array} \right), \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 3 & 1 & 2 \end{array} \right), \left( \begin{array}{rrrr} 1 & 2 & 3 \\ 2 & 1 & 3 \end{array} \right) \right\}$$

is not a subgroup of  $S_3$ .

- 30. Let G be any group and let g be a fixed element of G then prove, using the 2-step subgroup test, that  $H = \{gag^{-1} | a \in G\}$  is a subgroup.
- 31. Let G be any group then prove, using the 2-step subgroup test, that  $H = \{a \in G | ag = ga \forall g \in G\}$  is a subgroup.
- 32. For the groups  $G_1 = (\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2, +)$  and  $G_2 = (\mathbb{Z}_8, +)$ 
  - (a) Find the orders of the elements (1, 1, 1) and (1, 0, 1) in  $G_1$  and the orders of the elements 1, 2, 3 in  $G_2$ ?
  - (b) Are any of the above elements generators for their respective groups?
  - (c) Why aren't the groups isomorphic?
- 33. For the groups  $G_1 = (\mathbb{Z}_9^*, \cdot)$  and  $G_2 = (Z_6, +)$ 
  - (a) Find the orders of three different non identity elements in  $G_1$  and the orders of the elements 1, 2, 3 in  $G_2$ ?
  - (b) Are any of the above elements generators for their respective groups?
  - (c) The two groups are isomorphic. Find the isomorphism  $f: G_1 \to G_2$ .
- 34. Note that  $(G, \cdot)$  is a group where  $G = \{2^n : n \in \mathbb{Z}\}$  and  $\cdot$  is regular multiplication. Prove Axioms  $G_1$  and  $G_3$  for  $(G, \cdot)$ .
- 35. Note that  $(G, \cdot)$  is a group where  $G = \{2^n : n \in \mathbb{Z}\}$  and  $\cdot$  is regular multiplication. Show  $G \cong \mathbb{Z}$  where  $\mathbb{Z}$  is a group over addition. I used the isomorphism  $f : \mathbb{Z} \to G$ . We need to prove f preserves the operation and that f is a bijection.
- 36. Let  $(G_1, \cdot_1, e_1)$  and  $(G_2, \cdot_2, e_2)$  be groups and let  $f : G_1 \to G_2$  be an isomorphism. Show the following
  - (a) Then  $f(e_1) = e_2$
  - (b) Then  $f(a^{-1}) = f(a)^{-1}$