### Math 6250: Final Exam Review

#### Cardinality 1

- 1. What is the definition A and B have the same cardinality.
- 2. What is the definition A is countable.
- 3. List five sets which are countably infinite. List three sets that are uncountable.
- 4. Show the following sets have the same cardinality
  - (a)  $\mathbb{N}$  and  $\mathbb{Z}$
  - (b)  $\mathbb{N}$  and  $\mathbb{Q}$
  - (c) (1,5) and (3,11)

### $\mathbf{2}$ Some Complex Questions

- 5. Find all  $z \in \mathbb{C}$  so that
  - $z^3 = 1$
  - $z^3 = i$
  - $z^2 = i$
  - $z^4 = 1$
  - $z^2 = \frac{1}{2} + \frac{\sqrt{3}}{2}i$   $z^4 = 16$
- 6. Let  $z = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$ . Graph  $z, z^2, z^3$ . Describe what happens graphically when we square z. Look at the argument.
- 7. Let  $z = \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{2}}i$  and w = i. Graph z, w and zw. Describe what happens graphically when we multiply z and w. Look at the argument.
- 8. Let  $z \in \mathbb{C}$ . Prove  $z\overline{z}$  is a positive real number.
- 9. Use Euler's equation to prove a familiar trigonometric identity for

 $\cos(3\alpha)$ .

10. Use Euler's equation to prove a familiar trigonometric identity for

$$\sin(\alpha + \beta).$$

## 3 Limits of Sequences

11. Compute the following limits and use the  $\varepsilon - N$  definition to prove it.

(a) 
$$\lim_{n \to \infty} \frac{1}{3n^2 + 5n + 1}$$
  
(b) 
$$\lim_{n \to \infty} \frac{n}{3n + 1}$$
  
(c) 
$$\lim_{n \to \infty} \frac{3n + \sin(n)}{3 - 4n^2}$$

- 12. Use the  $\varepsilon N$  definition to prove If  $(a_n)$  is a convergent sequence then  $(a_n)$  is bounded.
- 13. Use the  $\varepsilon N$  definition to prove
  - (a) If  $\lim a_n = a$  and  $k \in \mathbb{R}$  then  $\lim ka_n = ka$
  - (b) If  $\lim a_n = a$  and  $\lim b_n = b$  then  $\lim a_n + b_n = a + b$
  - (c) If  $\lim a_n = a$  and  $\lim b_n = b$  then  $\lim a_n b_n = ab$

### 4 Limits of Functions

- 14. Compute the following limits and use the  $\varepsilon \delta$  definition to prove it.
  - (a)  $\lim_{x \to 3} x^2 + 2x$
  - (b)  $\lim_{x \to c} x^2 + 2x$
  - (c)  $\lim_{x\to -1} x^3 + 2x^2 + 1$  Hint divide  $x^3 + 2x^2 + 1$  by x + 1.
  - (d)  $\lim_{x\to 9}\sqrt{x}=3$
- 15. Use the  $\varepsilon \delta$  definition to prove
  - (a) If  $\lim_{x\to c} f(x) = F$  and  $k \in \mathbb{R}$  then  $_{x\to c} k f(x) = kF$
  - (b) If  $\lim_{x\to c} f(x) = F$  and  $\lim_{x\to c} g(x) = G$  then  $\lim_{x\to c} f(x) + g(x) = F + G$

16. Use the fact that

$$\lim_{a \to 0} \frac{\sin(a)}{a} = 1$$

to solve the following (do not use  $\varepsilon - \delta$ ):

(a)  $\lim_{x \to 0} \frac{1 - \cos(x)}{x}$ (b)  $\lim_{x \to 0} \frac{1 - \cos(x)}{x^2}$ (c)  $\lim_{x \to 0} \frac{\sin^2(x)}{x^2}$ (d)  $\lim_{h \to 0} \frac{\sin(x+h) - \sin(x)}{h}$ . Hint us Problem ??

## 5 Continuity

- 17. Show the following functions are continuus (or not).
  - (a)  $f(x) = x^{3}$  at x = c(b)  $f(x) = \frac{1}{x}$  at x = c(c)  $f(x) = \frac{x^{2} + x}{|x|}$  at x = 0(d)  $f(x) = \frac{x^{2} + x}{|x|}$  at x = 0(e)  $f(x) = \frac{x^{3} + x^{2}}{|x|}$  at x = 0(f)  $f(x) = \frac{x^{4} + x^{2}}{|x|}$  at x = 0(g)  $f(x) = \begin{cases} x \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$ (h)  $f(x) = \begin{cases} x \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$ (i)  $f(x) = \begin{cases} \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$ (j)  $f(x) = \begin{cases} x^{2} \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$ (k)  $f(x) = \begin{cases} x^{3} \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$

#### Derivatives 6

- 18. Prove if f is differentiable at x = c then f is continuous at x = c.
- 19. From the definition compute the derivatives for the following functions.

(a) 
$$f(x) = x^3$$
 at  $x = c$   
(b)  $f(x) = \frac{1}{x}$  at  $x = c$   
(c)  $f(x) = \frac{x^2 + x}{|x|}$  at  $x = 0$   
(d)  $f(x) = \frac{x^2 + x}{|x|}$  at  $x = 0$   
(e)  $f(x) = \frac{x^3 + x^2}{|x|}$  at  $x = 0$   
(f)  $f(x) = \frac{x^4 + x^2}{|x|}$  at  $x = 0$   
(g)  $f(x) = \begin{cases} x \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$   
(h)  $f(x) = \begin{cases} x \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$   
(i)  $f(x) = \begin{cases} \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$   
(j)  $f(x) = \begin{cases} x^2 \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$   
(k)  $f(x) = \begin{cases} x^3 \sin(\frac{1}{x}) & x \neq 0 \\ 0 & x = 0 \end{cases}$ 

20. Recall that

$$\lim_{\alpha \to 0} \frac{\sin(\alpha)}{\alpha} = 1.$$

Use this fact to prove the following

- (a)  $\lim_{\alpha \to 0} \frac{1 \cos(\alpha)}{\alpha} = 0.$ (a)  $\lim_{\alpha \to 0} \frac{\sin^2(\alpha)}{\alpha^2} = 1.$ (b)  $\lim_{\alpha \to 0} \frac{1 - \cos^2(\alpha)}{\alpha^2} = 1.$ (c)  $\lim_{\alpha \to 0} \frac{1 - \cos^2(\alpha)}{\alpha^2} = 0.$ (d)  $\lim_{\alpha \to 0} \frac{[1 - \cos(\alpha)]^2}{\alpha^2} = 0.$ (e)  $\lim_{\alpha \to 0} \frac{\tan(\alpha)}{\alpha} = 1.$

21. Note for  $f(x) = \sin(x)$  we have shown

$$\lim_{\alpha \to 0} \frac{1 - \cos(\alpha)}{\alpha} = 0.$$

And recall the trigonometric identity from Problem 9 Compute the derivative of f(x) using these two facts.

# 7 Trigonometry and Dimensions

- 22. Compute the Taylor series for the following functions at c = 0.
  - (a)  $f(x) = (x 3)^4$
  - (b)  $f(x) = \sin(2x)$
  - (c)  $f(x) = e^{5x}$