#### Math 3160 - Final Review

Be certain to know the Test 1 Review, Test 2 Review and . . .

### 1 Change of Basis Matrix

- 1. Let  $B = \{(1,0), (0,1)\}, B_1 = \{(-1,1), (2,3)\}$  and  $B_2 = \{(1,-1), (1,1)\}.$ 
  - (a) Find the change of basis matrices for  $P_{B_1 \to B_2}$  and  $P_{B_1 \to B_2}$ .
  - (b) Find the coordinates of the point (4,6) (given in the standard basis) relative to the bases  $B_1$  and  $B_2$ .
  - (c) Find the change of basis matrices for  $P_{B\to B_2}$  and  $P_{B_2\to B}$ .
  - (d) Find the coordinates of the point (2, -4) (given in the standard basis) relative to the bases B and  $B_2$ . Graph this point the two separate coordinate axes B and  $B_2$ .

### 2 Column Space & Null space

2. Let 
$$A = \begin{bmatrix} -1 & 2 & 0 & 3 & 0 \\ 2 & 1 & 1 & -1 & 1 \\ 1 & 3 & 1 & 2 & 1 \end{bmatrix}$$
.

- (a) Find a basis for the Column Space of A, COL(A).
- (b) Compute the dimension of COL(A).
- (c) Find a basis for the null space of A, NULL(A).
- (d) Compute the dimension of NULL(A).
- 3. The linear transformation  $T : \mathbb{R}^3 \to \mathbb{R}^3$  is given by the formula  $T(\begin{bmatrix} x \\ y \\ z \end{bmatrix}) = \begin{bmatrix} x+y \\ y+z \\ x-z \end{bmatrix}.$ 
  - (a) Find the matrix, A, to represent the linear transformation T.
  - (b) Compute the basis for the Range of T, which is the Column Space of A.
  - (c) Find a basis for the null space of A, NULL(A).

- (d) Compute the dimension of COL(A) and NULL(A). The dimension of the range of T is called the rank of T and the dimension of the null space is called the nullity.
- (e) What is the dimension of the domain of T and the codomain of T? Again, compare Rank, Nullity and the dimension of the Domain. Do you see a relation?

## **3** Diagonalizations

4. Find the eigenvalues and eigenvetors for the following matrices

$$A = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}, B = \begin{bmatrix} 3 & 4 \\ 2 & 3 \end{bmatrix}, C = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \text{ and } D = \begin{bmatrix} 4 & 0 & 0 \\ 3 & 2 & 1 \\ 0 & 4 & 0 \end{bmatrix}$$

5. Find the eigenvalues and eigenvetors for the following matrices (may have complex numbers in answer).

$$A = \begin{bmatrix} 1 & 1 \\ -1 & 3 \end{bmatrix}, \text{ and } B = \begin{bmatrix} 3 & 3 & 0 \\ 0 & 5 & 0 \\ 1 & 0 & 5 \end{bmatrix}.$$

6. Can the following matrices be diagonalized? State why? If it can be diagonalized, do it.

$$A = \begin{bmatrix} -1 & 2 \\ 2 & 1 \end{bmatrix}, C = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \text{ and } D = \begin{bmatrix} 4 & 0 & 0 \\ 3 & 2 & 1 \\ 0 & 4 & 0 \end{bmatrix}.$$

# 4 Cryptography - Hill Cipher

7. Which of the following can be used to encipher with the Hill cipher (as we did in class mod 26)?

$$A = \begin{bmatrix} 1 & 3 \\ 1 & 2 \end{bmatrix}, B = \begin{bmatrix} 4 & 1 \\ 1 & 2 \end{bmatrix}, C = \begin{bmatrix} 25 & 5 \\ 3 & 11 \end{bmatrix} \text{ and } D = \begin{bmatrix} 1 & 3 & 3 & 0 \\ 0 & 2 & 1 & 1 \\ 0 & 3 & 1 & 0 \\ 1 & 0 & 2 & 0 \end{bmatrix}$$

- 8. Define the following matrix:  $E = \begin{bmatrix} 5 & 1 \\ 3 & 6 \end{bmatrix}$ 
  - (a) For the plaintext message "WOLFMAN", find the two letter block representation using the alphabet:  $A = 00, B = 01, C = 02, D = 03, \ldots$
  - (b) Use the matrix E to encipher the plaintmessage.
  - (c) Find the deciphering matrix D.
  - (d) Decipher the ciphertext: JBCMVPTKGD.

## 5 Markov Chain

9. For the following defined matrices which are stochastic. 
$$A = \begin{bmatrix} .1 & .9 \\ .1 & .9 \end{bmatrix}$$
,  
 $B = \begin{bmatrix} .9 & 0 \\ .1 & 1 \end{bmatrix}$ ,  $C = \begin{bmatrix} 0.25 & -1.0 \\ 0.75 & 2.0 \end{bmatrix}$  and  $D = \begin{bmatrix} .1 & .2 & .3 & .4 \\ .0 & .4 & .1 & .5 \\ .0 & .4 & .6 & 0 \\ .9 & 0 & 0 & .1 \end{bmatrix}$ 

10. For the following defined markov processes which are regular.

$$A = \begin{bmatrix} .1 & .9 \\ .1 & .9 \end{bmatrix}, B = \begin{bmatrix} .9 & 0 \\ .1 & 1 \end{bmatrix} \text{ and } C = \begin{bmatrix} 0.25 & 1.0 \\ 0.75 & 0.0 \end{bmatrix}$$

- 11. Find the steady states for the following processes.
  - (a) For the transition matrix  $A = \begin{bmatrix} .1 & .9 \\ .9 & .1 \end{bmatrix}$ . You may be able to guess the answer for this one (but do the work anyway).
  - (b) A bike rental company has three locations. A renter can pick up a bike at any location and drop that bike off at any location.
    - A bike picked up at location A has 20% probability of being dropped off at location B and has a 10% probability of being dropped off at location C.
    - A bike picked up at location B has 30% probability of being dropped off at location A and has a 0% probability of being dropped off at location C.
    - A bike picked up at location C has 10% probability of being dropped off at location B and has a 40% probability of being dropped off at location A.