Math 3160 - Test 2

Name:

No calculators, no phones, no electronics allowed. Remember, I am grading your work. I must see how you came to the conclusion.

- 1. Let P(0,1,1), Q(0,1,0) and R(1,1,1) be points in \mathbb{R}^3 .
 - (a) Find the angle between the vectors \overrightarrow{PQ} and \overrightarrow{PR} .
 - (b) Compute the area of the parallelogram formed by the vectors \overrightarrow{PQ} and \overrightarrow{PR} .
 - (c) Find the normal vector of the plane containing P, Q and R.

2. Define the planes P_1 and P_2 as follows:

$$P_1: x - 2y + z = 12$$

$$P_2: y+z=4$$

- (a) Compute the angle between P_1 and P_2 .
- (b) What is the equation of the line of intersection of P_1 and P_2 ?

3. Let $\mathbf{v} = (1, 2, 0)$ and $\mathbf{w} = (0, -2, 1)$

- (a) Compute the projection of ${\bf v}$ onto ${\bf w}.$
- (b) Compute the parallel component of \mathbf{v} onto \mathbf{w} .
- (c) Compute the cooresponding perpendicular component of ${\bf v}$ onto ${\bf w}.$

4. Two step subspace test. Let $W = \{(a, b, c) \in \mathbb{R}^3 : a + 2b - c = 0\}$. Use the two step subspace test to show $(W, +, \cdot)$ is a subspace of \mathbb{R}^3 .

5. Let $B = \{(1,0,1), (-1,1,0), (2,2,0)\}$. Is the set of vectors independent? Show your work.

6. Let $B = \{(1,0,1), (-1,1,0), (0,2,2)\}$. Show the set of vectors is not independent, by finding a nontrivial linear combination of the vectors equal to the zero vector.

7. Let $B = \{(1,1), (-1,1)\}.$

- (a) Does B span \mathbb{R}^2 ? Show your work.
- (b) Show $(1,2) \in \text{span}(B)$ by finding a linear combination of the vectors in B equal to (1,2).

8. The linear transformation T is given by the formula

$$T\left(\left[\begin{array}{c} x\\y\\z\end{array}\right]\right) = \left[\begin{array}{c} x+y\\y+z\\x-z\\2x+y-z\end{array}\right].$$

- (a) What are the domain and codomain of T?
- (b) Find the matrix, A, to represent the linear transformation T.
- (c) Compute the basis for the column space of A, COL(A).
- (d) Compute the dimension of the COL(A). The dimension of the column space is called the rank.

9. The linear transformation T is given by the formula

$$T\left(\left[\begin{array}{c} x\\y\\z\end{array}\right]\right) = \left[\begin{array}{c} x+y\\y+z\\x-z\\2x+y-z\end{array}\right].$$

- (a) Compute the basis for the null space of A, NULL(A).
- (b) Compute the dimension of the NULL(A). The dimension of the null space is called the nullity.
- (c) What is the dimension of the domain of T and the codomain of T? What do you expect the dimension of the Range of T to be?

ec. Let $B = {\mathbf{v_1}, \mathbf{v_2}, \mathbf{v_3}}$ be a set of in a vector space, V.

- (a) Write the definition of Linear Independence of B.
- (b) Prove if there exists an $a_1, a_2 \in \mathbb{R} \setminus \{0\}$ so that

$$v_3 = a_1 \mathbf{v_1} + a_2 \mathbf{v_2}$$

then B is dependent.